

Powers in a class of \mathcal{A} -strict standard episturmian words

Amy Glen

`amy.glen@adelaide.edu.au`

`http://www.maths.adelaide.edu.au/~aglen`

The University of Adelaide



Introduction

Sturmian words



Introduction

Sturmian words

-  Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.




Introduction

Sturmian words

-  Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
-  Sturmian words are over a 2-letter alphabet.

Introduction

Sturmian words

-  Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
-  Sturmian words are over a 2-letter alphabet.
-  They are exactly the aperiodic infinite words of minimal complexity.

Introduction

● Sturmian words

- Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
- Sturmian words are over a 2-letter alphabet.
- They are exactly the aperiodic infinite words of minimal complexity.

● Episturmian words

Introduction

● Sturmian words

- Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
- Sturmian words are over a 2-letter alphabet.
- They are exactly the aperiodic infinite words of minimal complexity.

● Episturmian words

- A natural generalization of Sturmian words to an arbitrary finite alphabet.

Introduction

● Sturmian words

- Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
- Sturmian words are over a 2-letter alphabet.
- They are exactly the aperiodic infinite words of minimal complexity.

● Episturmian words

- A natural generalization of Sturmian words to an arbitrary finite alphabet.
- Introduced by Droubay, Justin, and Pirillo (2001).

Introduction

● Sturmian words

- Infinite words with $n + 1$ distinct factors of length n for each $n \in \mathbb{N}$.
- Sturmian words are over a 2-letter alphabet.
- They are exactly the aperiodic infinite words of minimal complexity.

● Episturmian words

- A natural generalization of Sturmian words to an arbitrary finite alphabet.
- Introduced by Droubay, Justin, and Pirillo (2001).
- Sturmian words are exactly the aperiodic episturmian words over a 2-letter alphabet.

Aim

- Explicitly determine all integer powers occurring in episturmian words.
- This has been done for Sturmian words by Damanik & Lenz (2003).
- We do this for a restricted class of episturmian words.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

● The *length* of u is $|u| = m$.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

- The *length* of u is $|u| = m$.
- The *reversal* of u is $\tilde{u} = x_mx_{m-1} \cdots x_1$.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

- The *length* of u is $|u| = m$.
- The *reversal* of u is $\tilde{u} = x_mx_{m-1} \cdots x_1$.
- u is a *palindrome* if $u = \tilde{u}$.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

- The *length* of u is $|u| = m$.
- The *reversal* of u is $\tilde{u} = x_mx_{m-1} \cdots x_1$.
- u is a *palindrome* if $u = \tilde{u}$.
- u^ω denotes the *purely periodic* infinite word $uuu \cdots$.

Terminology and notation

Let \mathcal{A} denote a *finite alphabet* and let $u = x_1x_2 \cdots x_m$ where each $x_i \in \mathcal{A}$.

- The *length* of u is $|u| = m$.
- The *reversal* of u is $\tilde{u} = x_mx_{m-1} \cdots x_1$.
- u is a *palindrome* if $u = \tilde{u}$.
- u^ω denotes the *purely periodic* infinite word $uuu \cdots$.
- For $0 \leq j \leq m-1$, the *j -th conjugate* of u is the word

$$C_j(u) := x_{j+1}x_{j+2} \cdots x_mx_1x_2 \cdots x_j$$

and we define

$$\mathcal{C}(u) := \{C_j(u) : 0 \leq j \leq |u| - 1\},$$

the *conjugacy class* of u .

Let x be an **infinite word** over \mathcal{A} .

● A *factor* of x is a finite string of consecutive letters in x .

Let x be an **infinite word** over \mathcal{A} .

- A *factor* of x is a finite string of consecutive letters in x .
- $\Omega(x)$ denotes the **set of all factors** of x .
- $\Omega_n(x)$ denotes the **set of all factors of x of length n** .

Let x be an **infinite word** over \mathcal{A} .

- A *factor* of x is a finite string of consecutive letters in x .
- $\Omega(x)$ denotes the **set of all factors** of x .
- $\Omega_n(x)$ denotes the **set of all factors of x of length n** .
- If $w \in \Omega(x)$, we write $w \prec x$.

Let x be an **infinite word** over \mathcal{A} .

- A *factor* of x is a finite string of consecutive letters in x .
- $\Omega(x)$ denotes the **set of all factors** of x .
- $\Omega_n(x)$ denotes the **set of all factors of x of length n** .
- If $w \in \Omega(x)$, we write $w \prec x$.
- A factor w of x is

$$\begin{cases} \text{right} \\ \text{left} \end{cases} \text{ *special* if } \begin{cases} wa, wb \\ aw, bw \end{cases}$$

are factors of x for some $a, b \in \mathcal{A}$, $a \neq b$.

Episturmian words

- An infinite word t is *episturmian* if:
 - $\Omega(t)$ is closed under reversal, and
 - t has at most one right special factor of each length.

Episturmian words

- An infinite word t is *episturmian* if:
 - $\Omega(t)$ is closed under reversal, and
 - t has at most one right special factor of each length.
- An episturmian word is *standard* if all of its left special factors are prefixes of it.

Standard episturmian words

- Let t be a standard episturmian word over \mathcal{A} and let

$$u_1 = \varepsilon, u_2, u_3, u_4, \dots$$

be the sequence of its palindromic prefixes.

Standard episturmian words

- Let t be a standard episturmian word over \mathcal{A} and let

$$u_1 = \varepsilon, u_2, u_3, u_4, \dots$$

be the sequence of its palindromic prefixes.

- \exists an infinite word $\Delta(t) = x_1 x_2 x_3 \cdots$ ($x_i \in \mathcal{A}$) such that

$$u_{n+1} = (u_n x_n)^{(+)}, \quad n \in \mathbb{N}^+.$$

Note: $w^{(+)}$ is the shortest palindrome of which w is a prefix.

Standard episturmian words

- Let t be a standard episturmian word over \mathcal{A} and let

$$u_1 = \varepsilon, \quad u_2, \quad u_3, \quad u_4, \quad \dots$$

be the sequence of its palindromic prefixes.

- \exists an infinite word $\Delta(t) = x_1x_2x_3 \cdots$ ($x_i \in \mathcal{A}$) such that

$$u_{n+1} = (u_n x_n)^{(+)}, \quad n \in \mathbb{N}^+.$$

Note: $w^{(+)}$ is the shortest palindrome of which w is a prefix.

- $\Delta(t)$ is called the *directive word* of t .

Standard episturmian words

- Let \mathbf{t} be a standard episturmian word over \mathcal{A} and let

$$u_1 = \varepsilon, \quad u_2, \quad u_3, \quad u_4, \quad \dots$$

be the sequence of its palindromic prefixes.

- \exists an infinite word $\Delta(\mathbf{t}) = x_1x_2x_3 \cdots$ ($x_i \in \mathcal{A}$) such that

$$u_{n+1} = (u_n x_n)^{(+)}, \quad n \in \mathbb{N}^+.$$

Note: $w^{(+)}$ is the shortest palindrome of which w is a prefix.

- $\Delta(\mathbf{t})$ is called the *directive word* of \mathbf{t} .
- $\mathbf{t} = \lim_{n \rightarrow \infty} u_n$

A class of strict standard episturmian words

Take $\mathcal{A} = \mathcal{A}_k := \{a_1, a_2, \dots, a_k\}$.

A class of strict standard episturmian words

Take $\mathcal{A} = \mathcal{A}_k := \{a_1, a_2, \dots, a_k\}$.

• t is *k -strict* if each $a_i \in \mathcal{A}_k$ appears infinitely often in $\Delta(t)$.

A class of strict standard episturmian words

Take $\mathcal{A} = \mathcal{A}_k := \{a_1, a_2, \dots, a_k\}$.

- t is *k -strict* if each $a_i \in \mathcal{A}_k$ appears infinitely often in $\Delta(t)$.
- k -strict episturmian words are precisely the k -letter Arnoux-Rauzy sequences.

A class of strict standard episturmian words

Take $\mathcal{A} = \mathcal{A}_k := \{a_1, a_2, \dots, a_k\}$.

- \mathbf{t} is *k -strict* if each $a_i \in \mathcal{A}_k$ appears infinitely often in $\Delta(\mathbf{t})$.
- k -strict episturmian words are precisely the k -letter **Arnoux-Rauzy sequences**.
- For any standard episturmian word \mathbf{t} ,

$$\Delta(\mathbf{t}) = a_1^{d_1} a_2^{d_2} \cdots a_k^{d_k} a_1^{d_{k+1}} a_2^{d_{k+2}} \cdots a_k^{d_{2k}} a_1^{d_{2k+1}} \cdots ,$$

where each $d_i \geq 0$.

A class of strict standard episturmian words

Take $\mathcal{A} = \mathcal{A}_k := \{a_1, a_2, \dots, a_k\}$.

- \mathbf{t} is *k -strict* if each $a_i \in \mathcal{A}_k$ appears infinitely often in $\Delta(\mathbf{t})$.
- k -strict episturmian words are precisely the k -letter **Arnoux-Rauzy sequences**.
- For any standard episturmian word \mathbf{t} ,

$$\Delta(\mathbf{t}) = a_1^{d_1} a_2^{d_2} \cdots a_k^{d_k} a_1^{d_{k+1}} a_2^{d_{k+2}} \cdots a_k^{d_{2k}} a_1^{d_{2k+1}} \cdots ,$$

where each $d_i \geq 0$.

- We restrict our attention to the case when all $d_i > 0$.
- Let \mathbf{s} be the **k -strict standard episturmian word** with directive word:

$$\Delta(\mathbf{s}) = a_1^{d_1} a_2^{d_2} \cdots a_k^{d_k} a_1^{d_{k+1}} a_2^{d_{k+2}} \cdots a_k^{d_{2k}} a_1^{d_{2k+1}} \cdots , \quad d_i > 0.$$

Example

- Let $\alpha \in (0, 1)$ be irrational with $\alpha = [0; 1 + d_1, d_2, d_3, \dots]$.
- The *characteristic Sturmian word* c_α over $\{a, b\}$ has directive word

$$\Delta(c_\alpha) = a^{d_1} b^{d_2} a^{d_3} b^{d_4} a^{d_5} \dots$$

Example

- Let $\alpha \in (0, 1)$ be irrational with $\alpha = [0; 1 + d_1, d_2, d_3, \dots]$.
- The *characteristic Sturmian word* c_α over $\{a, b\}$ has directive word

$$\Delta(c_\alpha) = a^{d_1} b^{d_2} a^{d_3} b^{d_4} a^{d_5} \dots$$

- $c_\alpha = \lim_{n \rightarrow \infty} s_n$, where $(s_n)_{n \geq -1}$ is defined by

$$s_{-1} = b, \quad s_0 = a, \quad s_n = s_{n-1}^{d_n} s_{n-2}, \quad n \geq 1.$$

Example

- Let $\alpha \in (0, 1)$ be irrational with $\alpha = [0; 1 + d_1, d_2, d_3, \dots]$.
- The *characteristic Sturmian word* c_α over $\{a, b\}$ has directive word

$$\Delta(c_\alpha) = a^{d_1} b^{d_2} a^{d_3} b^{d_4} a^{d_5} \dots$$

- $c_\alpha = \lim_{n \rightarrow \infty} s_n$, where $(s_n)_{n \geq -1}$ is defined by

$$s_{-1} = b, \quad s_0 = a, \quad s_n = s_{n-1}^{d_n} s_{n-2}, \quad n \geq 1.$$

- $\Delta(s)$ resembles $\Delta(c_\alpha)$.

Example

- Let $\alpha \in (0, 1)$ be irrational with $\alpha = [0; 1 + d_1, d_2, d_3, \dots]$.
- The *characteristic Sturmian word* c_α over $\{a, b\}$ has directive word

$$\Delta(c_\alpha) = a^{d_1} b^{d_2} a^{d_3} b^{d_4} a^{d_5} \dots$$

- $c_\alpha = \lim_{n \rightarrow \infty} s_n$, where $(s_n)_{n \geq -1}$ is defined by

$$s_{-1} = b, \quad s_0 = a, \quad s_n = s_{n-1}^{d_n} s_{n-2}, \quad n \geq 1.$$

- $\Delta(s)$ resembles $\Delta(c_\alpha)$.
- One can prove that $s = \lim_{n \rightarrow \infty} s_n$ where the sequence $(s_n)_{n \geq 1-k}$ is defined by

$$s_{1-k} = a_2, \quad s_{2-k} = a_3, \quad \dots, \quad s_{-1} = a_k, \quad s_0 = a_1,$$

$$s_n = s_{n-1}^{d_n} s_{n-2}^{d_{n-1}} \cdots s_0^{d_1} a_{n+1}, \quad 1 \leq n \leq k-1,$$

$$s_n = s_{n-1}^{d_n} s_{n-2}^{d_{n-1}} \cdots s_{n-k+1}^{d_{n-k+2}} s_{n-k}, \quad n \geq k.$$

Powers

- Let $p \geq 2$ be an integer.
- A finite word w has a *p -th power* in s if

$$w^p = \underbrace{ww \cdots w}_p \text{ is a factor of } s.$$

Powers

- Let $p \geq 2$ be an integer.
- A finite word w has a *p -th power* in s if

$$w^p = \underbrace{ww \cdots w}_p \text{ is a factor of } s.$$

Key tools in our analysis of powers occurring in s :

- canonical decompositions of s in terms of its **building blocks** s_n ;
- a generalization of **singular words**.

Singular words

- The set of factors of length $|s_n|$ in c_α is given by

$$\{\text{all conjugates of } s_n\} \cup \{w_n\}$$

where w_n is called the n -th *singular factor* of c_α .

[Wen and Wen (1994), Melançon (1999), Cao and Wen (2003)]

Singular words

- The set of factors of length $|s_n|$ in c_α is given by

$$\{\text{all conjugates of } s_n\} \cup \{w_n\}$$

where w_n is called the n -th *singular factor* of c_α .

[Wen and Wen (1994), Melançon (1999), Cao and Wen (2003)]

Singular n -words of the i -th kind

- The set of factors of s of length $|s_n|$ can be partitioned into k sets.

Singular words

- The set of factors of length $|s_n|$ in c_α is given by

$$\{\text{all conjugates of } s_n\} \cup \{w_n\}$$

where w_n is called the n -th *singular factor* of c_α .

[Wen and Wen (1994), Melançon (1999), Cao and Wen (2003)]

Singular n -words of the i -th kind

- The set of factors of s of length $|s_n|$ can be partitioned into k sets.

- That is:

$$\Omega_{|s_n|}(s) = \mathcal{C}(s_n) \dot{\cup} \Omega_n^1 \dot{\cup} \dots \dot{\cup} \Omega_n^{k-1}.$$

Singular words

- The set of factors of length $|s_n|$ in c_α is given by

$$\{\text{all conjugates of } s_n\} \cup \{w_n\}$$

where w_n is called the n -th *singular factor* of c_α .

[Wen and Wen (1994), Melançon (1999), Cao and Wen (2003)]

Singular n -words of the i -th kind

- The set of factors of s of length $|s_n|$ can be partitioned into k sets.

- That is:

$$\Omega_{|s_n|}(s) = \mathcal{C}(s_n) \dot{\cup} \Omega_n^1 \dot{\cup} \dots \dot{\cup} \Omega_n^{k-1}.$$

- Each set Ω_n^i is closed under reversal.

Singular words

- The set of factors of length $|s_n|$ in c_α is given by

$$\{\text{all conjugates of } s_n\} \cup \{w_n\}$$

where w_n is called the n -th *singular factor* of c_α .

[Wen and Wen (1994), Melançon (1999), Cao and Wen (2003)]

Singular n -words of the i -th kind

- The set of factors of s of length $|s_n|$ can be partitioned into k sets.

- That is:

$$\Omega_{|s_n|}(s) = \mathcal{C}(s_n) \dot{\cup} \Omega_n^1 \dot{\cup} \dots \dot{\cup} \Omega_n^{k-1}.$$

- Each set Ω_n^i is closed under reversal.
- If $w \in \Omega_n^i$ then w is called a *singular n -word of the i -th kind*.
- Such words play a key role in our study of powers occurring in s .

Powers occurring in s

- Let $n \in \mathbb{N}^+$ be fixed.
- We define k sets of lengths between $|s_n|$ and $|s_{n+1}|$:

$$\mathcal{D}_1(n) := \{r|s_n| : 1 \leq r \leq d_{n+1}\},$$

$$\mathcal{D}_i(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-i}^{d_{n+3-i}} s_{n+1-i}| : 1 \leq r \leq d_{n+1}\}, \quad 2 \leq i \leq k-1,$$

$$\mathcal{D}_k(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-k}^{d_{n+3-k}} s_{n+1-k}| : 1 \leq r \leq d_{n+1} - 1\}.$$

Powers occurring in s

- Let $n \in \mathbb{N}^+$ be fixed.
- We define k sets of lengths between $|s_n|$ and $|s_{n+1}|$:

$$\mathcal{D}_1(n) := \{r|s_n| : 1 \leq r \leq d_{n+1}\},$$

$$\mathcal{D}_i(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-i}^{d_{n+3-i}} s_{n+1-i}| : 1 \leq r \leq d_{n+1}\}, \quad 2 \leq i \leq k-1,$$

$$\mathcal{D}_k(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-k}^{d_{n+3-k}} s_{n+1-k}| : 1 \leq r \leq d_{n+1} - 1\}.$$

$$\text{Let } \mathcal{D}_n = \bigcup_{i=1}^k \mathcal{D}_i(n).$$

Powers occurring in s

- Let $n \in \mathbb{N}^+$ be fixed.
- We define k sets of lengths between $|s_n|$ and $|s_{n+1}|$:

$$\mathcal{D}_1(n) := \{r|s_n| : 1 \leq r \leq d_{n+1}\},$$

$$\mathcal{D}_i(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-i}^{d_{n+3-i}} s_{n+1-i}| : 1 \leq r \leq d_{n+1}\}, \quad 2 \leq i \leq k-1,$$

$$\mathcal{D}_k(n) := \{|s_n^r s_{n-1}^{d_n} \cdots s_{n+2-k}^{d_{n+3-k}} s_{n+1-k}| : 1 \leq r \leq d_{n+1} - 1\}.$$

Let $\mathcal{D}_n = \bigcup_{i=1}^k \mathcal{D}_i(n)$.

- Suppose $w \prec s$ and let $p \geq 2$ be an integer. Then,

$$w^p \prec s \quad \Rightarrow \quad |w| \in \mathcal{D}_n \text{ for some } n.$$

Squares

- In s , successive occurrences of a singular word are *positively separated*.

Squares

- In s , successive occurrences of a singular word are *positively separated*.

- Consequently:

Lemma: Suppose $u^2 \prec s$ with $|u| \in \mathcal{D}_n$. Then

$$w \not\prec u \quad \text{if} \quad w \in \Omega_{n+1-i}^1 \text{ for some } i \in [1, k-1].$$

Squares

- In s , successive occurrences of a singular word are *positively separated*.

- Consequently:

Lemma: Suppose $u^2 \prec s$ with $|u| \in \mathcal{D}_n$. Then

$$w \not\prec u \quad \text{if} \quad w \in \Omega_{n+1-i}^1 \text{ for some } i \in [1, k-1].$$

- That is, u does not contain a singular $(n+1-i)$ -word of the first kind for any $i \in [1, k-1]$.

Squares, cubes, and higher powers

• Let $w \prec s$ with $|w| \in \mathcal{D}_n$ for some n .

Squares, cubes, and higher powers

● Let $w \prec s$ with $|w| \in \mathcal{D}_n$ for some n .

● Our main results show:

If $w^p \prec s$, then w is a conjugate of a finite product of blocks from the set $\{s_n, s_{n-1}, \dots, s_{n+1-k}\}$, depending on $|w|$ and d_{n+1} .

Squares, cubes, and higher powers

● Let $w \prec s$ with $|w| \in \mathcal{D}_n$ for some n .

● Our main results show:

If $w^p \prec s$, then w is a conjugate of a finite product of blocks from the set $\{s_n, s_{n-1}, \dots, s_{n+1-k}\}$, depending on $|w|$ and d_{n+1} .

● For instance:

● Let $p \geq 2$.

● Suppose $|w| = r|s_n|$ for some r with $1 \leq r < (d_{n+1} + 2)/p$.

Squares, cubes, and higher powers

● Let $w \prec s$ with $|w| \in \mathcal{D}_n$ for some n .

● Our main results show:

If $w^p \prec s$, then w is a conjugate of a finite product of blocks from the set $\{s_n, s_{n-1}, \dots, s_{n+1-k}\}$, depending on $|w|$ and d_{n+1} .

● For instance:

● Let $p \geq 2$.

● Suppose $|w| = r|s_n|$ for some r with $1 \leq r < (d_{n+1} + 2)/p$.

● Then:

$$w^p \prec s \iff w \text{ is one of the first } |s_n| \text{ conjugates of } (s_n)^r.$$

Example: k -bonacci word

- Define the *k -bonacci word* to be the standard episturmian word η_k with directive word $(a_1 a_2 \cdots a_k)^\omega$.

Example: k -bonacci word

- Define the *k -bonacci word* to be the standard episturmian word η_k with directive word $(a_1 a_2 \cdots a_k)^\omega$.
- Since all $d_i = 1$, we have $s_n = s_{n-1} s_{n-2} \cdots s_{n-k}$ for all $n \geq 1$.
(The lengths $|s_n|$ are the *k -bonacci numbers*.)

Example: k -bonacci word

- Define the *k -bonacci word* to be the standard episturmian word η_k with directive word $(a_1 a_2 \cdots a_k)^\omega$.
- Since all $d_i = 1$, we have $s_n = s_{n-1} s_{n-2} \cdots s_{n-k}$ for all $n \geq 1$.
(The lengths $|s_n|$ are the *k -bonacci numbers*.)
- If $w^p \prec \eta_k$, then
$$|w| = |s_n| + |s_{n-1}| + \cdots + |s_{n+1-i}| \quad \text{for some } n \in \mathbb{N} \text{ and } i \in [1, k-1].$$

-
- Our main results reveal that, in η_k ,
 - $(a_1)^2$ is the unique square of length 2;
 - all conjugates of s_n have a square;
 - only some conjugates of s_n have a cube;
 - only some conjugates of $s_n s_{n-1} \cdots s_{n+1-i}$ have a square.

-
- Our main results reveal that, in η_k ,
 - $(a_1)^2$ is the unique square of length 2;
 - all conjugates of s_n have a square;
 - only some conjugates of s_n have a cube;
 - only some conjugates of $s_n s_{n-1} \cdots s_{n+1-i}$ have a square.
 - There are no other integer powers in η_k .
 - In particular, the k -bonacci word is 4-power free.

Concluding remarks

- Our main results on powers suffice to describe all integer powers occurring in any (episturmian) word that is equivalent to s .

- **Open problem:**

Determine all integer powers occurring in general standard episturmian words (with not all d_i necessarily positive).